

WELSH JOINT EDUCATION COMMITTEE CYD-BWYLLGOR ADDYSG CYMRU

General Certificate of Education

Tystysgrif Addysg Gyffredinol

Advanced Level/Advanced Subsidiary

Safon Uwch/Uwch Gyfrannol

MATHEMATICS FP3

Further Pure Mathematics

Specimen Paper 2005/2006

(1½ hours)

INSTRUCTIONS TO CANDIDATES

Answer **all** questions.

INFORMATION FOR CANDIDATES

A calculator may be used for this paper.

A formula booklet is available and may be used.

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

1. Solve the equation

$$\cosh^2 x = 3 + \sinh x,$$

expressing the roots as natural logarithms.

[7]

2. (a) By drawing appropriate graphs, show that the equation

$$x^3 = \cot x$$

has one root in the interval $(0, \pi/2)$.

[3]

- (b) Starting with an initial approximation $x_0 = 1$, use the Newton-Raphson method to calculate successive approximations x_1 , x_2 and x_3 to this root. Write down the value of x_3 correct to 6 decimal places and determine whether or not this gives the value of the root correct to 6 decimal places.

[8]

3. The arc joining the points $(0,0)$ and $(1,1)$ on the curve $y = x^3$ is rotated through four right-angles about the x -axis.

- (a) (i) Show that the area of the curved surface generated is given by

$$2\pi \int_0^1 x^3 \sqrt{1+9x^4} dx.$$

[2]

- (ii) Use the substitution $u = 1 + 9x^4$ to show this area is equal to

$$\frac{\pi}{27}(10\sqrt{10} - 1).$$

[6]

4. Given that

$$I = \int_0^{\pi/2} e^{-2x} \cos x dx$$

$$\text{and } J = \int_0^{\pi/2} e^{-2x} \sin x dx,$$

use integration by parts to show that

$$I = e^{-\pi} + 2J$$

$$\text{and } J = 1 - 2I$$

Hence evaluate I and J , giving each answer in the form $a + be^{-\pi}$, where a and b are rational numbers.

[11]

5. (a) Find the Maclaurin series of $\ln(1 + \sin x)$ up to and including the x^3 term. [9]

- (b) Use your series to evaluate, approximately, the integral

$$\int_0^{\frac{1}{3}} \ln(1 + \sin x) dx. \quad [4]$$

6. The curves C_1 and C_2 have polar equations as follows:

$$C_1 : r = 1 - \cos \theta \quad (-\pi \leq \theta \leq \pi)$$

$$C_2 : r = \cos 2\theta \quad \left(-\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}\right)$$

- (a) Sketch C_1 and C_2 on the same diagram. [2]
- (b) Find the area enclosed by C_1 . [5]
- (c) Find the polar coordinates of the points of intersection of C_1 and C_2 . [6]

7. (a) Show that

$$\frac{\sin n\theta - \sin(n-1)\theta}{\sin \theta} = \cos(n-1)\theta. \quad [2]$$

- (b) Given that

$$I_n = \int_0^\pi \frac{\sin n\theta}{\sin \theta} d\theta$$

where n is an integer, show that for $n \geq 2$,

$$I_n = I_{n-2}. \quad [4]$$

- (c) Hence evaluate I_n when n is

- (i) an even integer,
- (ii) an odd integer. [6]